

Manuscript Draft

Manuscript Number: AMC-D-07-02659

Title: Generalized FLT

Article Type: Short Communication

Keywords: FTT, Primes, Co primes

Corresponding Author: Mr hemant ajay pandey,

Corresponding Author's Institution: City Montessary school

First Author: hemant ajay pandey

Order of Authors: hemant ajay pandey

Abstract: The Abstract: The generalized Fermat's last theorem is a open problem in number theory and it asks whether the equation

$xa + yb = zc$ has a solution $x, y, z, l + 1$ & $a, b, c, 3$ and x, y, z are mutually co-prime.

As we all are well aware of the celebrated proof of Fermat's last theorem by legendry Sir Andrew wiles in 1974. The truth of FLT in general case when all the exponents are different integers larger than 3, is the main point of discussion of present paper.

We have attempted to establish the truth of the result through the method of contradiction.

Our method from the start assumes that x, y, z , are mutually co prime and in the progress of our solution we try to establish that for a solution they can't be co prime.

1

2 *Title Page*

3 1. *Contents*

4 2. *Author s Profile*

5 3. *Cover Letter*

6 4. *Abstract*

7 5. *Manuscript*

8 6. *References*

9

10

11

12

13

14

15

16

17

18

19

AUTHORS PROFILE

Author: Hemant Pandey

AffiliationPresently working as an IIT JEE lecturer for TIME institute:

City Montessary School (CMS) –Innovation Wing (IW),Amelox college tutor

www.amelox.com

Job Profile: Content Developer; Writer,IIT JEE lecturer

Branch: Main Branch,

Station Road, Lucknow

Writing profile: Currently working on a text book for

Grade 11-12 of mathematics with Amelox College Tutor; California (USA)

Qualification: Masters of Science in Mathematics and currently a Researcher

Postal Address:

16/732, Indira Nagar,Lucknow.

Uttar Pradesh – 226016(Pin) India

Mobile Number +91-9336291008, +91-9338202560

Mailing Address:

E-mail- ajay_fermat@rediffmail.com

46 E-mail- ajay_euler@rediffmail.com

47 E-mail- ajay_gauss@rediffmail.com

48

49

50

51

52

53

Cover Letter

54

55 *Dear Sir,*

56 *My name is hemant pandey and I am a young researcher.*

57 *I am proposing an paper on generalized FLT for possible publication. The*
58 *main argument used in the paper is the method of contradiction and*
59 *property of all mutually co prime numbers, that each number in the universe*
60 *is expressible as a linear combination of two co prime numbers.*

61 *This is to certify that the work is original and has not been submitted any*
62 *where else for publication consideration.*

63 *Details are in the pdf attachment.*

64

65 *Sincerely,*

66 *Hemant Pandey*

67

68

69

70

71

A Generalization to FLT

The Abstract: The generalized Fermat's last theorem is a open problem in number theory and it asks whether the equation

$x^a + y^b = z^c$ has a solution $\forall x, y, z, \in I^+$ & $a, b, c, \geq 3$ and x, y, z are mutually co-prime.

As we all are well aware of the celebrated proof of Fermat's last theorem by legendry Sir Andrew wiles in 1974. The truth of FLT in general case when all the exponents are different integers larger than 3, is the main point of discussion of present paper.

We have attempted to establish the truth of the result through the method of contradiction.

Our method from the start assumes that x, y, z , are mutually co prime and in the progress of our solution we try to establish that for a solution they can't be co prime.

97

98

99

100

101 ***Meaning of symbols:***

102 x^k - *x raised to the power 'k'*

103 \forall - *For all*

104 I^+ - *set of + ve integers*

105 \Rightarrow - *Implies that*

106 $>$ - *Is greater than*

107 $<$ - *Is less than*

108 $=$ - *Is equal to*

109 \neq - *Is not equal to*

110 \in - *Belongs to*

111

112

113

114

115

116

117

The Manuscript

1. Introduction

The generalization of FLT can be expressed in a classic one line statement:

The following equation has no solution, for all $x, y, z \in I^+$ and mutually co-prime.

a, b, c are integers greater than 2.

$\therefore x^a + y^b \neq z^c \dots[A]$

If x, y, z are co prime, $\in I^+$

And a, b, c are integers greater than or equal to three.

The following proposed proof works on the method of contradiction.

2.Proof:

We would start our proof with a known result that for all x, y, z mutually co-prime we can find l, m , mutually co-prime

s.t $lx-my = 1$ note that l & y and x & m are coprime.

$\Rightarrow z = (l \cdot z) \cdot x - (m \cdot z) \cdot y$

140 $\forall z \in I^+$

141 \Rightarrow We can always find $A, B \forall x, y, z$, where the three variables are co prime.

142 s.t $z = Ax - By$, where $A=l.z$ and $B=m.z$

143 Rewriting (A) we get

144 $x^a + y^b \neq (Ax - By)^c$

145 is equivalent generalized Fermat's last theorem.

146 \therefore To employ the method of contradiction assuming the contrary

147 Let $x^a + y^b = (Ax - By)^c$ for some $\forall x, y$ co prime (B)

148 Note back since $A = l.z$

149 $B = m.z$

150

151 A, y & B, x , are mutually co prime as y & z and x & z are mutually co prime.

152 (R-1)

153 Now (B) gives

154 $x^a + y^b = (A.x)^c - (A.x)^{c-1} \cdot {}^cC_1 \cdot (By) + \dots + (-1)^c \cdot (By)^c$

155 Rearranging we get $x^a - (A.x)^c = -y^b - (Ax)^{c-1} \cdot {}^cC_1 \cdot (By) + \dots + (-1)^c \cdot (By)^c$

156 $x^a[1 - A^c(x)^{c-a}] = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-1} \cdot (By)^{c-1} \}$

157 or $x^a[1 - A^c(x)^{c-a}] = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot (B^c \cdot y^{c-1}) \} \dots \dots \dots (C)$

158 Since x & y co-prime

159 $\Rightarrow 1 - (A.x)^{c-a} = \lambda_1 y$ [where $\lambda_1 = \text{integer}$](1)

160 Now putting this value in (C) we have

161

162 $x^a \lambda_1 y = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot (B^c \cdot y^{c-1}) \}$

163 Or $x^a \lambda_1 = y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot B^c \cdot y^{c-1}$

164 $\Rightarrow x^a \lambda_1 = y^{b-1} + x[A^{c-1} \cdot x^{c-2} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} A] + (-1)^{c-1} \cdot B^c \cdot y^{c-1}$

165 Or $x^a \lambda_1 = y^{b-1} + (-1)^{c-1} \cdot B^c \cdot y^{c-1} + x[A^{c-1} \cdot x^{c-2} - {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot A]$

166

167 Or $x^a \lambda_1 = y^{b-1} [1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b}] + x [A^{c-1} \cdot x^{c-2} - {}^c C_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot {}^2 A]$

168

169 $\Rightarrow y^{b-1} + (-1)^{c-1} \cdot B^c \cdot y^{c-1} = \lambda_2 x$

170 Since other summand is a multiple of x .

171 Or $y^{b-1} \{1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b}\} = \lambda_2 x$

172

173 Now x & y are co prime

174 $\Rightarrow 1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b} = \lambda_2 x \dots (2)$

175 from 1 & 2 we have

176 $1 - (A)^c x^{c-a} = \lambda_1 y$ & $1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b} = \lambda_2 x \dots (3)$

177 Here λ_1 & λ_2 are arbitrary integers to be determined.

178 Note from above situation $a \neq b \neq c$

179 if $a = b = c$ the conditions cannot hold good, as the variables become unity.

180 This excludes the cases of FLT to be encountered here.

181 $a \neq b \neq c$ is also a condition.

182 Here we have tacitly assumed that $c > a > b$ but that does not affect generality.

183 As if $a > b > c$ we may take

184 $x^a = z^c - y^b$ etc.

185 all the other steps are similar

186 The greatest exponent is to be equated to the sum or difference of other two.

187 Now set (3) gives that

188 $x(A)^c \cdot x^{c-a-1} + \lambda_1 y = 1$

189 & $\lambda_2 \cdot x + y(B^c \cdot y^{c-b-1})(-1)^c = 1$

190 A little algebra gives that for a solution

$$191 \quad \frac{A^{c-a} \cdot x^{c-a-1}}{\lambda_2} = \frac{\lambda_1}{B^c \cdot y^{c-b} \cdot (-1)^c} = 1$$

$$192 \quad \lambda_2 = A^c \cdot x^{c-a-1}$$

$$193 \quad \lambda_1 = B^c \cdot y^{c-b-1} \cdot (-1)^c$$

194 *Putting the above values in the equation (C) we get*

$$195 \quad x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c = -y^b [1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b-1}] - y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

$$196 \quad {}^2 \cdot x \cdot A \cdot y^c$$

$$197 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c = (-1) \{y^b \cdot A^c \cdot x^{c-a-1} \cdot x\} - y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

$$198 \quad {}^2 \cdot x \cdot A \cdot y^c$$

$$199 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c + \{y^b \cdot A^c \cdot x^{c-a}\} = -y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

200

$$201 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c + \{y^b \cdot A^c \cdot x^{c-a}\} + (A \cdot x)^c + (-1)^c (By)^c$$

$$202 \quad = -y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A] + (A \cdot x)^c + (-1)^c (By)^c$$

$$203 \quad \Rightarrow x^a \cdot [B^c \cdot y^{c-b} \cdot (-1)^c + A^c \cdot x^{c-a}] + y^b \cdot [A^c \cdot x^{c-a} + B^c \cdot y^{c-b} \cdot (-1)^c] = (Ax - By)^c$$

204

205 *Comparing with (B) we have*

206

$$207 \quad B^c \cdot y^{c-b} \cdot (-1)^c + A^c \cdot x^{c-a} = 1$$

208 *Which is certainly impossible, since B=l.z & A=m.z*

209 *Only possibility may be z=1, But that case is taken care of via catalan's conjecture's*
 210 *proof.*

211 *We may formally prove the case even.*

212

$$213 \quad Z=1 \Rightarrow m^c \cdot y^{c-b} \cdot (-1)^c + l^c \cdot x^{c-a} = 1$$

$$214 \quad \text{But } l \cdot x - m \cdot y = 1$$

$$215 \quad \Rightarrow l^c \cdot x^{c-a-1} / l = m^c \cdot y^{c-b-1} \cdot (-1)^c / -m$$

$$216 \quad \Rightarrow l^{c-1} \cdot x^{c-a-1} = m^{c-1} \cdot y^{c-b-1} \cdot (-1)^{c-1}$$

217

218 *Which is certainly untrue since l , y & m are co prime etc.*

219 *Hence $x^a + y^b = z^c$*

220 *has no soln. \forall*

221 *$a \# b \# c$*

222 *& x, y, z co prime.*

223 *Hence the result.*

224

225

226

227 Sole Author

228 Hemant Pandey

229

1

2 *Title Page*

3 1. *Contents*

4 2. *Author s Profile*

5 3. *Cover Letter*

6 4. *Abstract*

7 5. *Manuscript*

8 6. *References*

9

10

11

12

13

14

15

16

17

18

19

AUTHORS PROFILE

Author: Hemant Pandey

AffiliationPresently working as an IIT JEE lecturer for TIME institute:

City Montessary School (CMS) –Innovation Wing (IW),Amelox college tutor

www.amelox.com

Job Profile: Content Developer; Writer,IIT JEE lecturer

Branch: Main Branch,

Station Road, Lucknow

Writing profile: Currently working on a text book for

Grade 11-12 of mathematics with Amelox College Tutor; California (USA)

Qualification: Masters of Science in Mathematics and currently a Researcher

Postal Address:

16/732, Indira Nagar,Lucknow.

Uttar Pradesh – 226016(Pin) India

Mobile Number +91-9336291008, +91-9338202560

Mailing Address:

E-mail- ajay_fermat@rediffmail.com

46 E-mail- ajay_euler@rediffmail.com

47 E-mail- ajay_gauss@rediffmail.com

48

49

50

51

52

53

Cover Letter

54

55 *Dear Sir,*

56 *My name is hemant pandey and I am a young researcher.*

57 *I am proposing an paper on generalized FLT for possible publication. The*
58 *main argument used in the paper is the method of contradiction and*
59 *property of all mutually co prime numbers, that each number in the universe*
60 *is expressible as a linear combination of two co prime numbers.*

61 *This is to certify that the work is original and has not been submitted any*
62 *where else for publication consideration.*

63 *Details are in the pdf attachment.*

64

65 *Sincerely,*

66 *Hemant Pandey*

67

68

69

70

71

A Generalization to FLT

The Abstract: The generalized Fermat's last theorem is a open problem in number theory and it asks whether the equation

$x^a + y^b = z^c$ has a solution $\forall x, y, z, \in I^+$ & $a, b, c, \geq 3$ and x, y, z are mutually co-prime.

As we all are well aware of the celebrated proof of Fermat's last theorem by legendry Sir Andrew wiles in 1974. The truth of FLT in general case when all the exponents are different integers larger than 3, is the main point of discussion of present paper.

We have attempted to establish the truth of the result through the method of contradiction.

Our method from the start assumes that x, y, z , are mutually co prime and in the progress of our solution we try to establish that for a solution they can't be co prime.

97

98

99

100

101 ***Meaning of symbols:***

102 x^k - *x raised to the power 'k'*

103 \forall - *For all*

104 I^+ - *set of + ve integers*

105 \Rightarrow - *Implies that*

106 $>$ - *Is greater than*

107 $<$ - *Is less than*

108 $=$ - *Is equal to*

109 \neq - *Is not equal to*

110 \in - *Belongs to*

111

112

113

114

115

116

117

The Manuscript

1. Introduction

The generalization of FLT can be expressed in a classic one line statement:

The following equation has no solution, for all $x, y, z \in I^+$ and mutually co-prime.

a, b, c are integers greater than 2.

$\therefore x^a + y^b \neq z^c \dots[A]$

If x, y, z are co prime, $\in I^+$

And a, b, c are integers greater than or equal to three.

The following proposed proof works on the method of contradiction.

2.Proof:

We would start our proof with a known result that for all x, y, z mutually co-prime we can find l, m , mutually co-prime

s.t $lx-my = 1$ note that l & y and x & m are coprime.

$\Rightarrow z = (l \cdot z) \cdot x - (m \cdot z) \cdot y$

140 $\forall z \in I^+$

141 \Rightarrow We can always find $A, B \forall x, y, z$, where the three variables are co prime.

142 s.t $z = Ax - By$, where $A=l.z$ and $B=m.z$

143 Rewriting (A) we get

144 $x^a + y^b \neq (Ax - By)^c$

145 is equivalent generalized Fermat's last theorem.

146 \therefore To employ the method of contradiction assuming the contrary

147 Let $x^a + y^b = (Ax - By)^c$ for some $\forall x, y$ co prime (B)

148 Note back since $A = l.z$

149 $B = m.z$

150

151 A, y & B, x , are mutually co prime as y & z and x & z are mutually co prime.

152 (R-1)

153 Now (B) gives

154 $x^a + y^b = (A.x)^c - (A.x)^{c-1} \cdot {}^cC_1 \cdot (By) + \dots + (-1)^c \cdot (By)^c$

155 Rearranging we get $x^a - (A.x)^c = -y^b - (Ax)^{c-1} \cdot {}^cC_1 \cdot (By) + \dots + (-1)^c \cdot (By)^c$

156 $x^a[1 - A^c(x)^{c-a}] = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-1} \cdot (By)^{c-1} \}$

157 or $x^a[1 - A^c(x)^{c-a}] = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot (B^c \cdot y^{c-1}) \} \dots \dots \dots (C)$

158 Since x & y co-prime

159 $\Rightarrow 1 - (A.x)^{c-a} = \lambda_1 y$ [where $\lambda_1 = \text{integer}$](1)

160 Now putting this value in (C) we have

161

162 $x^a \lambda_1 y = -y \{ y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot (B^c \cdot y^{c-1}) \}$

163 Or $x^a \lambda_1 = y^{b-1} + (A.x)^{c-1} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-1} \cdot B^c \cdot y^{c-1}$

164 $\Rightarrow x^a \lambda_1 = y^{b-1} + x[A^{c-1} \cdot x^{c-2} \cdot {}^cC_1 \cdot B - \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} A] + (-1)^{c-1} \cdot B^c \cdot y^{c-1}$

165 Or $x^a \lambda_1 = y^{b-1} + (-1)^{c-1} \cdot B^c \cdot y^{c-1} + x[A^{c-1} \cdot x^{c-2} - {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot A]$

166

167 Or $x^a \lambda_1 = y^{b-1} [1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b}] + x [A^{c-1} \cdot x^{c-2} - {}^c C_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot {}^2 A]$

168

169 $\Rightarrow y^{b-1} + (-1)^{c-1} \cdot B^c \cdot y^{c-1} = \lambda_2 x$

170 Since other summand is a multiple of x .

171 Or $y^{b-1} \{1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b}\} = \lambda_2 x$

172

173 Now x & y are co prime

174 $\Rightarrow 1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b} = \lambda_2 x \dots (2)$

175 from 1 & 2 we have

176 $1 - (A)^c x^{c-a} = \lambda_1 y$ & $1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b} = \lambda_2 x \dots (3)$

177 Here λ_1 & λ_2 are arbitrary integers to be determined.

178 Note from above situation $a \neq b \neq c$

179 if $a = b = c$ the conditions cannot hold good, as the variables become unity.

180 This excludes the cases of FLT to be encountered here.

181 $a \neq b \neq c$ is also a condition.

182 Here we have tacitly assumed that $c > a > b$ but that does not affect generality.

183 As if $a > b > c$ we may take

184 $x^a = z^c - y^b$ etc.

185 all the other steps are similar

186 The greatest exponent is to be equated to the sum or difference of other two.

187 Now set (3) gives that

188 $x(A)^c \cdot x^{c-a-1} + \lambda_1 y = 1$

189 & $\lambda_2 \cdot x + y(B^c \cdot y^{c-b-1})(-1)^c = 1$

190 A little algebra gives that for a solution

$$191 \quad \frac{A^{c-a} \cdot x^{c-a-1}}{\lambda_2} = \frac{\lambda_1}{B^c \cdot y^{c-b} \cdot (-1)^c} = 1$$

$$192 \quad \lambda_2 = A^c \cdot x^{c-a-1}$$

$$193 \quad \lambda_1 = B^c \cdot y^{c-b-1} \cdot (-1)^c$$

194 *Putting the above values in the equation (C) we get*

$$195 \quad x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c = -y^b [1 + (-1)^{c-1} \cdot B^c \cdot y^{c-b-1}] - y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

$$196 \quad {}^2 \cdot x \cdot A \cdot \}]$$

$$197 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c = (-1) \{ y^b \cdot A^c \cdot x^{c-a-1} \cdot x \} - y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

$$198 \quad {}^2 \cdot x \cdot A \cdot \}]$$

$$199 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c + \{ y^b \cdot A^c \cdot x^{c-a} \} = -y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A]$$

200

$$201 \quad \text{Or } x^a \cdot B^c \cdot y^{c-b} \cdot (-1)^c + \{ y^b \cdot A^c \cdot x^{c-a} \} + (A \cdot x)^c + (-1)^c (By)^c$$

$$202 \quad = -y [(Ax)^{c-1} \cdot {}^cC_1 \cdot B + \dots + (-1)^{c-2} \cdot B^{c-1} \cdot y^{c-2} \cdot x \cdot A] + (A \cdot x)^c + (-1)^c (By)^c$$

$$203 \quad \Rightarrow x^a \cdot [B^c \cdot y^{c-b} \cdot (-1)^c + A^c \cdot x^{c-a}] + y^b \cdot [A^c \cdot x^{c-a} + B^c \cdot y^{c-b} \cdot (-1)^c] = (Ax - By)^c$$

204

205 *Comparing with (B) we have*

206

$$207 \quad B^c \cdot y^{c-b} \cdot (-1)^c + A^c \cdot x^{c-a} = 1$$

208 *Which is certainly impossible, since B=l.z & A=m.z*

209 *Only possibility may be z=1, But that case is taken care of via catalan's conjecture's*
 210 *proof.*

211 *We may formally prove the case even.*

212

$$213 \quad Z=1 \Rightarrow m^c \cdot y^{c-b} \cdot (-1)^c + l^c \cdot x^{c-a} = 1$$

$$214 \quad \text{But } l \cdot x - m \cdot y = 1$$

$$215 \quad \Rightarrow l^c \cdot x^{c-a-1} / l = m^c \cdot y^{c-b-1} \cdot (-1)^c / -m$$

$$216 \quad \Rightarrow l^{c-1} \cdot x^{c-a-1} = m^{c-1} \cdot y^{c-b-1} \cdot (-1)^{c-1}$$

217

218 *Which is certainly untrue since l , y & m are co prime etc.*

219 *Hence $x^a + y^b = z^c$*

220 *has no soln. \forall*

221 *$a \# b \# c$*

222 *& x, y, z co prime.*

223 *Hence the result.*

224

225

226

227 Sole Author

228 Hemant Pandey

229